


TEST NAME: Cycle Mapping Plus

Estrogen (E) patterns can be seen below in green. Progesterone (Pg) patterns can be seen below in purple. Normal ranges are within the gray dashed lines. See page 2 for more information.



All values given in ng/mg creatinine

| Measurement | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 |
|-----------------|---|----|----|----|----|----|----|----|----|
| Day(s) of Cycle | 4 | 10 | 12 | 13 | 15 | 17 | 20 | 24 | 28 |

The days listed above were used for measurements. Two samples are used and listed for long cycles or patients without a normal cycle.

| | | | | | | | | | |
|-------------------------|------|------|------|------|------|------|------|------|------|
| Estradiol (E2) | 1.00 | 1.56 | 3.86 | 8.33 | 4.53 | 3.03 | 3.33 | 3.10 | 0.91 |
| Estrone (E1) | 3.1 | 3.9 | 6.8 | 27.5 | 16.2 | 13.2 | 12.0 | 12.7 | 4.8 |
| a-Pregnanediol | 57 | 35 | 41 | 42 | 45 | 115 | 179 | 230 | 86 |
| b-Pregnanediol | 162 | 127 | 138 | 147 | 157 | 312 | 467 | 565 | 256 |
| b-Pregnanediol/E2 Ratio | 161 | 81 | 36 | 18 | 35 | 103 | 140 | 182 | 282 |
| Creatinine | 1.00 | 1.75 | 1.90 | 1.65 | 2.41 | 1.23 | 1.11 | 0.87 | 0.86 |

Sample (#8) with the highest b-Pg value (565) is used for E and Pg metabolites for DUTCH Complete or Plus if ordered.


| Normal Ranges | Follicular | Ovulatory | Luteal | Postmenopausal |
|----------------|--------------|--------------|---------------|----------------|
| Estradiol | 1-2ng/mg | 4-12ng/mg | 1.8-4.5ng/mg | 0.2-0.7ng/mg |
| Estrone | 4-12ng/mg | 22-68ng/mg | 12-26ng/mg | 3.0-7.0ng/mg |
| a-Pregnanediol | 25-100ng/mg | 25-100ng/mg | 200-740ng/mg | 15-50ng/mg |
| b-Pregnanediol | 100-300ng/mg | 100-300ng/mg | 600-2000ng/mg | 60-200ng/mg |

b-Pregnanediol/E2 ratio is typically 50-300 in the follicular phase, <100 during ovulation, and 100-500 in the luteal phase. Creatinine normal range, 0.2-2.0 mg/mL. Values outside this range may be less certain due to under or overhydration.

TEST NAME: Cycle Mapping Plus

The comments below include general information that we hope you will find useful in your understanding of the patient's results. These results and comments are not intended to diagnose any specific conditions.

You'll find four stacked graphs with the reference ranges and the patient's results mapped out. The top graphs represent estrogen (E) production, and the bottom graphs represents progesterone (Pg) production. The horizontal axis shows the cycle days (0-30+) and the vertical axis shows hormone concentration or hormone metabolites being measured. Healthy cycles typically range from 21-35 days. The patient likely submitted many samples over one cycle, and we have selected the 9 most relevant measurements. Some measurements from longer cycles are from two-day averages to ensure transitory E and Pg peaks are not missed.




For most cycles <34 days, measurements are made from single days, selected to best represent overall patterns of ovulatory & luteal peaks. The day 4 sample set is usually collected at the end of the cycle, four days after menses (used for DUTCH Complete/Plus), but is plotted at the beginning of the cycle as above. If a DUTCH Complete or Plus was ordered, data for the E and Pg metabolite values are taken from the day on the Cycle Mapping associated with the progesterone (b-pregnane-diol) peak in the luteal phase.

The first part of the cycle (days 1-14) is the "follicular phase," ovulation typically occurs mid-cycle, and the "luteal phase" refers to the 2nd half of the cycle (days 14 until menses). These phases may shift in patients with atypical cycle lengths. Levels may still be considered normal in short or long cycles even if the timing of the E or Pg peaks are at different times.

In the top graphs, we follow both primary estrogens, estrone (E1) and estradiol (E2). In a typical cycle, estrogen rises in the follicular phase, which stimulates the luteinizing hormone (LH) surge from the brain about 24-36 hours before ovulation, which leads to the production of Pg the second half of the cycle (measured by its primary pregnane-diol metabolites). Pg rises only after ovulation has occurred, reaching its peak 5-7 days later, then begins to decrease before the onset of menses. When Pg does not rise it indicates that the patient is likely not ovulating. A weak rise in Pg can also indicate either no ovulation or a weak corpus luteum (luteal phase defect), which is associated with poor egg maturation, difficulty maintaining a secretory endometrium and infertility. Ranges for Pg are similar for a postmenopausal woman or a cycling woman who is in the follicular phase. In the table near the bottom of page 1 below the graphs, the patient's results are displayed in a table. This includes creatinine, which is used to correct for hydration. If creatinine is very low or very high, hormone measurements from that day may be less reliable.


Below are four different cycle patterns that may help with interpretation





TEST NAME: Cycle Mapping Plus

Hormone Testing Summary


Key (how to read the results):




Sex Hormones See Pages 2 and 3 for a thorough breakdown of sex hormone metabolites



Progesterone Serum Equivalent is a calculated value based on urine pregnanediol.





Adrenal Hormones See pages 4 and 5 for a more complete breakdown of adrenal hormones




Free cortisol best reflects tissue levels. Metabolized cortisol best reflects total cortisol production.

Total DHEA Production

| Age | Range |
|-------|-----------|
| 20-39 | 1300-3000 |
| 40-59 | 750-2000 |
| >60 | 500-1200 |



cortisol metabolism



PLEASE BE SURE TO READ BELOW FOR ANY SPECIFIC LAB COMMENTS. More detailed comments can be found on page 8.

- This DUTCH Test is associated with a Cycle Mapping report as well. The progesterone and estrogen metabolites for this report are based on the Cycle Mapping sample with the highest progesterone value. 2-Methoxy-E2 and 4-OH-E2 are not included for Cycle Mapping patients. The Cortisol Awakening Response (CAR) was 3.21ng/mL (expected range 1.5-4.0) or 315% (range 50-160%). See page 5 for more details.

TEST NAME: Cycle Mapping Plus

| Test | | Result | Units | Luteal* | Postmenopausal |
|--|--------------------------|--------|-------|------------|----------------|
| Progesterone Metabolites (Urine) | | | | | |
| b-Pregnanediol | Low end of luteal range | 611.0 | ng/mg | 600 - 2000 | 60-200 |
| a-Pregnanediol | High end of luteal range | 641.0 | ng/mg | 200 - 740 | 15-50 |
| Estrogens and Metabolites (Urine) | | | | | |
| Estrone(E1) | Within luteal range | 17.1 | ng/mg | 12 - 26 | 3.0-7.0 |
| Estradiol(E2) | Above luteal range | 5.34 | ng/mg | 1.8 - 4.5 | 0.2-0.7 |
| Estriol(E3) | Below luteal range | 2.9 | ng/mg | 5 - 18 | 0.6-4.0 |
| 4-OH-E1 | Within luteal range | 1.06 | ng/mg | 0 - 1.8 | 0-0.3 |
| 16-OH-E1 | Below luteal range | 0.48 | ng/mg | 0.7 - 2.6 | 0.2-0.6 |
| 2-Methoxy-E1 | | 5.58 | ng/mg | 2.5 - 6.5 | 0.3-1.4 |
| 2-OH-E2 | Within luteal range | 0.69 | ng/mg | 0 - 1.2 | 0-0.3 |
| Total Estrogen | Low end of range | 40.0 | ng/mg | 35 - 70 | 4.0-15 |


| Androgens and Metabolites (Urine) | | | | | |
|--|-------------------|--------|-------|------------|--|
| DHEA-S | High end of range | 637.0 | ng/mg | 20 - 750 | |
| Androsterone | Above range | 1996.0 | ng/mg | 200 - 1650 | |
| Etiocholanolone | High end of range | 847.0 | ng/mg | 200 - 1000 | |
| Testosterone | Below range | 1.3 | ng/mg | 2.3 - 14 | |
| 5a-DHT | Within range | 3.3 | ng/mg | 0 - 6.6 | |
| 5a-Androstanediol | Above range | 34.3 | ng/mg | 12 - 30 | |
| 5b-Androstanediol | Low end of range | 28.5 | ng/mg | 20 - 75 | |
| Epi-Testosterone | Low end of range | 2.6 | ng/mg | 2.3 - 14 | |

*the Luteal Range is the premenopausal range. When patients are taking oral progesterone this range for progesterone metabolites is not luteal and reflects the higher levels expected when patients take oral progesterone. This test is intended to be taken in the luteal phase of the menstrual cycle (days 19-22 of a 28 day cycle) for premenopausal women. The ranges in the table below may be used when samples are taken during the first few days (follicular) of the cycle, during ovulation (days 11-14) or when patients are on oral progesterone. See the following pages for age-dependent ranges for androgen metabolites.

| Additional Normal Ranges | Follicular | Ovulatory | Oral Pg (100mg) |
|--------------------------|------------|-----------|-----------------|
| b-Pregnanediol | 100-300 | 100-300 | 2000-9000 |
| a-Pregnanediol | 25-100 | 25-100 | 580-3000 |
| Estrone (E1) | 4.0-12.0 | 22-68 | N/A |
| Estradiol (E2) | 1.0-2.0 | 4.0-12.0 | N/A |

TEST NAME: Cycle Mapping Plus


Hormone metabolite results from the previous page are presented here as they are found in the steroid cascade. See the Provider Comments for more information on how to read the results.



TEST NAME: Cycle Mapping Plus

| Category | Test | Result | Units | Normal Range |
|--|--------------------------------|-------------------|--------|-------------------|
| Free Cortisol and Cortisone (Saliva) | | | | |
| | Saliva Cortisol - Waking (W) | Below range | 1.02 | ng/mL 1.6 - 4.6 |
| | Saliva Cortisol - W+30 min. | Low end of range | 4.23 | ng/mL 3.7 - 8.2 |
| | Saliva Cortisol - W+60 min. | Below range | 1.89 | ng/mL 2.3 - 5.3 |
| | Saliva Cortisol - Afternoon | Within range | 0.91 | ng/mL 0.4 - 1.5 |
| | Saliva Cortisol - Night | Low end of range | 0.04 | ng/mL 0 - 0.9 |
| | Saliva Cortisone - Waking (W) | Low end of range | 7.99 | ng/mL 6.8 - 14.5 |
| | Saliva Cortisone - W+30 min. | Within range | 14.1 | ng/mL 12.4 - 19.4 |
| | Saliva Cortisone - W+60 min. | Within range | 10.92 | ng/mL 9.4 - 15.3 |
| | Saliva Cortisone - Afternoon | Within range | 5.67 | ng/mL 2 - 7.1 |
| | Saliva Cortisone - Night | Within range | 1.65 | ng/mL 0 - 4.8 |
| | Saliva Cortisol Total | Below range | 8.08 | ng/mL 9.6 - 19.3 |
| | Saliva Cortisone Total | Within range | 40.33 | ng/mL 36 - 55 |
| Creatinine (Urine) | | | | |
| | Creatinine A (Waking) | Within range | 0.83 | mg/ml 0.2 - 2 |
| | Creatinine B (Morning) | Within range | 0.45 | mg/ml 0.2 - 2 |
| | Creatinine C (Afternoon) | Within range | 0.27 | mg/ml 0.2 - 2 |
| | Creatinine D (Night) | Within range | 0.78 | mg/ml 0.2 - 2 |
| Cortisol Metabolites and DHEA-S (Urine) | | | | |
| | a-Tetrahydrocortisol (a-THF) | Within range | 273.0 | ng/mg 75 - 370 |
| | b-Tetrahydrocortisol (b-THF) | Low end of range | 1335.0 | ng/mg 1050 - 2500 |
| | b-Tetrahydrocortisone (b-THE) | Within range | 2901.0 | ng/mg 1550 - 3800 |
| | Metabolized Cortisol (THF+THE) | Within range | 4509.0 | ng/mg 2750 - 6500 |
| | DHEA-S | High end of range | 637.0 | ng/mg 20 - 750 |

TEST NAME: Cycle Mapping Plus



The Cortisol Awakening Response (CAR) is the rise in salivary cortisol between the waking sample and the sample collected 30 (as well as 60) minutes later. This "awakening response" is essentially a "mini stress test" and is a useful measurement in addition to the overall up-and-down (diurnal) pattern of free cortisol throughout the day. **This patient shows a waking cortisol of 1.02 and an increase to 4.23 after 30.0 minutes. This is an increase of 3.21ng/mL or 315%.** Expected increases differ depending on the methods used. Preliminary research shows that 50-160% or 1.5-4.0ng/mL increases are common with samples collected 30 minutes after waking. These guidelines are considered research only. **This patient shows a salivary cortisol of 1.89 measured 60 minutes after waking. This is an increase of 0.87ng/mL or 85.3% compared to the waking sampe.** To date, data suggests that expected results may be 0-70%, and this guideline is considered for research only.



PATIENT: **XXXXXXXXXXXXXXXXXXXX**

TEST REF: **TST-NL-XXXX**

TEST NUMBER: T-NL-XXXXX (XXXXXXXXXXXX)

COLLECTED: XX/XX/XXXX

PRACTITIONER:

GENDER: XYZ

RECEIVED: XX/XX/XXXX

XXXXXXXXXXXXXXXXXXXX

AGE: XX

TESTED: XX/XX/XXXX

XXXXXXXXXXXXXXXXXXXX

TEST NAME: Cycle Mapping Plus

| Category | Test | Result | Units | Normal Range |
|--|----------------------|------------------|------------|--------------|
| Nutritional Organic Acids | | | | |
| Vitamin B12 Marker (may be deficient if high) - (Urine) | | | | |
| | Methylmalonate (MMA) | Within range | 0.7 ug/mg | 0 - 2.5 |
| Vitamin B6 Markers (may be deficient if high) - (Urine) | | | | |
| | Xanthurenate | Within range | 0.66 ug/mg | 0.12 - 1.2 |
| | Kynurenate | Within range | 2.4 ug/mg | 0.8 - 4.5 |
| Biotin Marker (may be deficient if high) - (Urine) | | | | |
| | b-Hydroxyisovalerate | Within range | 1.5 ug/mg | 0 - 12.5 |
| Glutathione Marker (may be deficient if low or high) - (Urine) | | | | |
| | Pyroglutamate | Below range | 25.3 ug/mg | 28 - 58 |
| Gut Marker (potential gut putrefaction or dysbiosis if high) - (Urine) | | | | |
| | Indican | Within range | 2.0 ug/mg | 0 - 100 |
| Neuro-related Markers | | | | |
| Dopamine Metabolite - (Urine) | | | | |
| | Homovanillate (HVA) | Low end of range | 3.3 ug/mg | 3 - 11 |
| Norepinephrine/Epinephrine Metabolite - (Urine) | | | | |
| | Vanilmandelate (VMA) | Low end of range | 2.2 ug/mg | 2.2 - 5.5 |
| Neuroinflammation Marker - (Urine) | | | | |
| | Quinolate | Within range | 5.0 ug/mg | 0 - 9.6 |
| Additional Markers | | | | |
| Melatonin (*measured as 6-OH-Melatonin-Sulfate) - (Urine) | | | | |
| | Melatonin* (Waking) | Low end of range | 13.9 ng/mg | 10 - 85 |
| Oxidative Stress / DNA Damage, measured as 8-Hydroxy-2-deoxyguanosine (8-OHdG) - (Urine) | | | | |
| | 8-OHdG (Waking) | Within range | 1.6 ng/mg | 0 - 5.2 |

TEST NAME: Cycle Mapping Plus




Provider Notes

How to read the DUTCH report


This report is not intended to treat, cure or diagnose any specific diseases. The graphic dutch dials in this report are intended for quick and easy evaluation of which hormones are out of range. Results below the left star are shaded yellow and are below range (left). Results between the stars and shaded green are within the reference range (middle). Results beyond the second star and shaded red are above the reference range (right). Some of these hormones also change with age, and the age-dependent ranges provided should also be considered.



For female reproductive hormones, a purple band is present on the dutch dials. This band represents the expected levels (reference range) for postmenopausal (or non-cycling) women.



In a few places on the graphical pages, you will see fan-style gauges. For sex hormones, you will see one for the balance between 5a/5b metabolism as well as methylation. For adrenal hormones, you will see one to represent the balance between cortisol and cortisone metabolites. These indexes simply look at the ratio of hormones for a preference. An average or "normal" ratio between the two metabolites (or groups of metabolites) will give a result in the middle (as shown here). If the ratio between the metabolites measured is "low" the gauge will lean to the left and similarly to the right if the ratio is higher than normal.



Patient or Sample Comments

Throughout the provider comments you may find some comments specific to your situation or results. These comments will be found in this section or within another section as appropriate. Comments in other sections that are specific to your case will be in **bold**.

The patient reports regular menstrual cycles.

The patient reported symptoms of excess estrogen. This can be caused by excess estrogen or progesterone deficiency. Results should be carefully reviewed. We do not report a progesterone to estrogen ratio. However, you can investigate this issue by looking at the relative level of these two hormones on their respective gauges.

The patient reported significant symptoms of excess androgen levels.

The patient is experiencing significant fatigue in the mornings.

Progesterone Metabolism

The primary role of progesterone is to balance the strong effects of estrogen. Progesterone metabolites are measured and reflect progesterone levels well because very little progesterone is found in urine, so b-Pregnanediol is typically used as a surrogate marker because it is the most abundant metabolite, but we also test the corresponding a-pregnanediol. The average of the two metabolites is reported for progesterone. If levels are in the lower part of the reference range compared to estrogen levels, symptoms of too much estrogen may occur.

When ordering the DUTCH Complete, you will see Progesterone Serum Equivalent on the summary page 1. The urine metabolites of progesterone have been proven to correlate strongly enough to serum progesterone to provide this value. The correlation is the strongest for values within the premenopausal luteal range. Urine metabolites can at times result in somewhat higher serum equivalent results in the postmenopausal range. For this reason the postmenopausal Serum Equivalent range is slightly higher than typical serum ranges. NOTE: If progesterone is taken orally (also with sublingual), these metabolites are elevated from gut metabolism and results do NOT accurately reflect serum levels.

Progesterone results are in range for a cycling woman in the luteal phase, indicating ovulation likely occurred 5-7 days before the sample collection. Evaluate if healthy progesterone levels are in balance with estrogen, especially if cyclical symptoms are a part of the picture.



PATIENT: XXXXXXXXXXXXXXXXXXXX

TEST REF: TST-NL-XXXX

TEST NUMBER: T-NL-XXXXX (XXXXXXXXXX)

COLLECTED: XX/XX/XXXX

PRACTITIONER:

GENDER: XYZ

RECEIVED: XX/XX/XXXX

XXXXXXXXXXXXXXXXXX

AGE: XX

TESTED: XX/XX/XXXX

XXXXXXXXXXXXXXXXXXXXXXXXXX

TEST NAME: Cycle Mapping Plus

As she metabolizes her progesterone, you can see she shows a preference for making the a-Pregnanediol. This metabolite can cross the blood-brain-barrier and works on GABA receptors, giving a calming effect. Women who metabolize their progesterone through this pathway tend to have more significant benefit from oral progesterone for symptoms of anxiety and difficulty sleeping.

Estrogen Metabolism

When evaluating estrogen levels, it is important to assess the following:

- **The status (low, normal or high?) of estrogen production:**

Levels of the primary ovarian product, estradiol (the strongest estrogen), as well as "total estrogens" may be considered. For women not on HRT, consider the appropriate range (premenopausal or postmenopausal).

- **Phase I Metabolism:**

Estrogen is metabolized (primarily by the liver) down three phase I pathways. The 2-OH pathway is considered the safest because of the anti-cancer properties of 2-OH metabolites. Conversely, the 4-OH pathway is considered the most genotoxic as its metabolites can create reactive products that damage DNA. The third pathway, 16-OH creates the most estrogenic of the metabolites (although still considerably less estrogenic than estradiol) - 16-OH-E1. If overall estrogen levels are high, production of 16-OH-E1 may exacerbate high estrogen symptoms. Similarly, a woman with very low levels of estrogens, may have less low estrogen symptoms if 16-OH metabolism is preferred. For example Armamento-Villareal showed that a higher 2-OH-E1/16-OH-E1 ratio correlated to bone loss (a low estrogen symptom). Estriol is thought of as a safer (weaker) estrogen metabolite, but it is important to remember that estriol is actually 16-OH-E2, so generally patients that make a lot of the potentially protective/weak estriol may also make a lot of the estrogenic 16-OH-E1.

When evaluating phase I metabolism, it may be important to look at the ratios of the three metabolites to see which pathways are preferred relative to one another. It may also be important to compare these metabolites to the levels of the parent hormones (E1, E2). If the ratios of the three metabolites are favorable but overall levels of metabolites are much lower than E1 and E2, this may imply sluggish phase I clearance of estrogens, which can contribute to high levels of E1 and E2. Similarly, patients with excessive phase I metabolism may have low E1 and E2 levels because of high rates of clearance (as opposed to simply not making a lot of estrogen).

The pie chart will assist you in comparing the three pathway options of phase I metabolism compared to what is "normal." 2-OH metabolism can be increased by using products containing D.I.M. or I-3-C. These compounds are found (or created from) in cruciferous vegetables and are known for promoting this pathway.

Phase I metabolism shows a preference for 2-OH metabolism, which is the protective pathway. The 4-OH and 16-OH pathways may be out of range despite this preference, so careful assessment is necessary. Products to increase 2-OH metabolism may be considered if E1 and E2 are elevated or if the 4-OH or 16-OH estrogens are high relative to 2-OH estrogens. Products that push the 2-OH pathway may also lower E1 and E2 levels, so keep this in mind when considering therapy.

- **Methylation (part of phase II metabolism) of estrogens:**

After phase I metabolism, both 4-OH and 2-OH (not 16-OH) estrogens can be deactivated and eliminated by methylation. The methylation-activity index shows the patient's ratio of 2-Methoxy-E1 / 2-OH-E1 compared to what is expected. Low methylation can be caused by low levels of nutrients needed for methylation and/or genetic abnormalities (COMT, MTHFR). The COMT enzyme responsible for methylation requires magnesium and methyl donors. Deficiencies in folate or vitamin B6 or B12 can cause low levels of methyl donors. MTHFR genetic defects can make it more difficult for patients to make sufficient methyl donors. Genetic defects in COMT can make methylation poor even in the presence of adequate methyl donors.

Androgen Metabolism

When evaluating androgen levels, it is important to assess the following:

- **The status (low, normal or high?) of DHEA:**

DHEA and androstenedione are made almost exclusively by the adrenal gland (although a smaller amount is made in the ovaries). These hormones appear in urine as DHEA-S (DHEA-Sulfate), androsterone and etiocholanolone. The best way to assess the total production of DHEA is to add up these three metabolites. This total can be seen on the first page of the DUTCH Complete (and DUTCH Plus). DHEA production decreases quite significantly with age. Age-dependent ranges can be seen on the graphical page of results.

The Total DHEA Production (page 1) was about 3,480ng/mg which is elevated. High DHEA can cause symptoms of androgen excess including oily skin, acne, sleep problems, headaches and mood disturbances. High levels may be due to supplementation, insulin, stress, elevated prolactin, alcohol and certain medications like ADD meds, Xanax and Wellbutrin. High DHEA can be treated with blood sugar balancing lifestyle, stress reduction and in appropriate cases ashwagandha. In some cases, highly androgenic people may show high levels of both DHEA or testosterone without negative clinical consequence.

- **The status (low, normal or high?) of testosterone:**

Females make most of their DHEA in the adrenal gland and a fraction of that DHEA trickles down metabolically to testosterone. For premenopausal women, some testosterone is also made by the ovaries. Levels of testosterone do drop somewhat with age, but not to the degree that DHEA decreases.

The patient noted being of Asian descent. There is a very common genetic variant in this population that, if present, disallows much of testosterone from being converted into the water-soluble form found in urine. Actual testosterone levels, if this genetic variant are present, would be better assessed in a blood test (preferably "total" and "free" testosterone). This variant metabolism has no known clinical impact (it is neither

TEST NAME: Cycle Mapping Plus


good nor bad but simply makes urine testosterone falsely lower than it would otherwise be) and impacts testosterone, 5a-DHT and 5b-androstanediol. It is not known to impact epi-testosterone, 5a-androstanediol or other androgens as they are metabolized (conjugated to their water-soluble forms) by a different enzyme.

Testosterone levels for this patient were approximately 1.30ng/mg, which is low. This can cause symptoms of low energy, low motivation, loss of lean muscle mass and low libido. Depending on her age, consider testosterone HRT or supplements that support low testosterone symptoms include Zinc, Maca, Tribulus, and Shatavari. If the patient does not present with symptoms of low testosterone, you may want to also carefully evaluate 5a-metabolism (see below) and testosterone's downstream metabolites, 5a-androstanediol and 5b-androstanediol to confirm a low androgenic state.

The metabolic preference for the 5a (5-alpha) or 5b (5-beta) pathway:

5a-reductase converts testosterone into 5a-DHT (DHT), which is even more potent (~3x) than testosterone. High levels of DHT can lead to symptoms associated with too much testosterone. Metabolites created down the 5b-pathway are significantly less androgenic than their 5a counterparts. In the examples below, the example on the left shows a patient with 5b-metabolism preference. A patient with a pattern like the example on the right may have high androgen symptoms even though the hormones are in the normal range because of the likely preference for turning a lot of her testosterone into DHT. The fan-style gauge below the hormones shows the 5a or 5b preference based on etiocholanolone (5b) and androsterone (5a) results. Progesterone metabolites are also metabolized by 5a and 5b enzymes and the balance between its two metabolites can be useful to confirm a 5a or 5b preference.

Example of how to read fan-style gauge for 5a-reductase activity:



While testosterone levels are not high, overall DHEA production is on the higher side and androgens are preferring the androgenic 5a pathway. Since the patient did list significant symptoms of high androgens, these higher levels may not be well tolerated by the patient. High insulin levels can lead to more DHEA production and 5a-metabolism, it may be worth exploring potential issues with blood sugar and/or insulin or possibly strategies to block 5a-metabolism (saw palmetto, zinc, nettles, pygeum and reishi). Some providers may also prefer to use pharmaceutical options (spironolactone or metformin) as it fits.

It is important to consider DHEA and testosterone production, 5a-metabolism patterns as well as the patient symptoms. For example, a woman with higher levels of DHEA and testosterone will often have high androgen symptoms (facial hair, thinning scalp hair, etc.) exacerbated by 5a-metabolism. If, on the other hand, she prefers 5b-metabolism she may not express high androgen symptoms in spite of higher levels of testosterone because 5b is the less androgenic pathway. Testosterone levels may be better understood by also considering its downstream metabolites (5a-androstanediol, 5b-androstanediol). Technically, these metabolites can also be formed from DHEA metabolites without going through the testosterone pathway, but they generally tend to correlate with testosterone production. You will also see levels of epi-testosterone, which is not androgenic like testosterone. It happens to be produced in about the same concentrations as testosterone (this is an approximate relationship). This can be helpful to assess testosterone therapy and rare cases where testosterone may have other complexities.

DUTCH Adrenal

The HPA-Axis refers to the communication and interaction between the hypothalamus (H) and pituitary (P) in the brain down to the adrenal glands (A) that sit on top of your kidneys. When a physical or psychological stressor occurs, the hypothalamus tells the pituitary to make ACTH, a hormone. ACTH stimulates the adrenal glands to make the stress hormone, cortisol and to a lesser extent DHEA and DHEA-S. Normally, the HPA-axis production follows a daily pattern in which cortisol rises rather rapidly in the first 10-30 minutes after waking (this is the C.A.R.) in order to help with energy, then gradually decreases throughout the day so that it is low at night for sleep. The cycle starts over the next morning. Abnormally high activity occurs in Cushing's Disease where the HPA-axis is hyper-stimulated causing cortisol to be elevated all day. The opposite is known as Addison's Disease, where cortisol is abnormally low because it is not made appropriately in response to ACTH's stimulation. These two conditions are somewhat rare. Examples of more common conditions related to less severely abnormal cortisol



| | | |
|--------------------------------------|-----------------------|--------------------------------|
| PATIENT: XXXXXXXXXXXXXXXXXXXX | TEST REF: TST-NL-XXXX | |
| TEST NUMBER: T-NL-XXXXX (XXXXXXXXXX) | COLLECTED: XX/XX/XXXX | PRACTITIONER: XXXXXXXXXXXXXXXX |
| GENDER: XYZ | RECEIVED: XX/XX/XXXX | XXXXXXXXXXXXXXXXXXXXX |
| AGE: XX | TESTED: XX/XX/XXXX | |

TEST NAME: Cycle Mapping Plus

levels include fatigue, depression, insomnia, fibromyalgia, anxiety, inflammation and more.

Only a fraction of cortisol is "free" and bioactive. This fraction of cortisol is very important, but levels of metabolized cortisol best represent overall production of cortisol therefore both should be taken into account to correctly assess adrenal function.

When evaluating cortisol levels, it is important to assess the following:

- **The overall up-and-down pattern of free cortisol throughout the day, looking for low and high levels:**

Abnormal results should be considered along with related symptoms.

- **The sum of the free cortisol as an expression of the overall tissue cortisol exposure:**

This total of five free cortisol measurements is the best way to assess the total of free cortisol throughout the day, but do be aware that it is heavily weighted towards the morning production since three of five measurements are made within the first hour of the day.

- **The total level of cortisol metabolites:**

We call this calculation "Metabolized Cortisol" which is the sum of a-THF, b-THF and b-THE. While free cortisol is the best assessment for tissue levels of cortisol, it only represents 1-3% of the total produced. The majority of cortisol results in a urine metabolite and the total of these metabolites best represents the total glandular output of cortisol for the day. When overall production is much higher than free cortisol levels, cortisol clearance may be increased (as seen in hyperthyroidism, obesity, etc.) The most common reason for sluggish cortisol clearance (assumed when free cortisol levels are much higher than metabolized cortisol) is low thyroid.

- **A potential preference for cortisol or cortisone (the inactive form):**

Looking at the comparison between the total for free cortisol and free cortisone is NOT the best indication of a person's preference for cortisol or cortisone. The saliva gland converts cortisol to cortisone in the local tissue. This localized conversion can be seen by comparing cortisol and cortisone levels. To see the patient's preference systemically, it is best to look at which *metabolite* predominates (THF or THE). This preference can be seen in the gauge below metabolized cortisol. This is known as the 11b-HSD index. The enzyme 11b-HSD II converts cortisol to cortisone in the kidneys, saliva gland and colon. 11b-HSD I is more active in the liver, fat cells and the periphery and is responsible for reactivating cortisone to cortisol. Both are then metabolized by 5a-reductase to become tetrahydrocortisol (THF) and tetrahydrocortisone (THE) respectively.

- **The Cortisol Awakening Response (CAR):**

The unique feature of the DUTCH Plus is the inclusion of the CAR assessment. The response to waking adds one more piece to HPA-axis function. In some cases overall levels of free cortisol may be normal, but the response to stress may be under or overactive. Reasons for a lower CAR might include: an underactive HPA Axis, excessive psychological burnout, seasonal affective disorder (SAD), sleep apnea or poor sleep in general, PTSD, and "chronic fatigue" patients. An elevated CAR can be a result of an over-reactive HPA axis, ongoing job-related stress (anticipatory stress for the day), glycemic dysregulation, pain (ie. waking with painful joints or a migraine), and general depression (not SAD). Scientific literature points to the magnitude of the morning cortisol increase as being connected to HPA-axis health whether the overall production of cortisol is low, normal or high.

Nutritional Organic Acids

The following three organic acids are functional markers for vitamin deficiency. These compounds essentially back up in human biochemistry when a key nutrient is missing. These three metabolites have fairly straightforward interpretations. When the markers are elevated, it is likely that the patient's cellular levels of the related nutrient may be insufficient.

Methylmalonate (MMA)

Methylmalonate (also known as methylmalonic acid or MMA) is a functional marker of vitamin B12 (also known as cobalamin) deficiency. When cellular levels of B12 are low either from deficiency or due to a B12 transporter gene mutation, levels of MMA increase. This marker is considered superior to measuring serum B12 levels directly. A 2012 publication by Miller showed that 20% of those tested had a genetic defect in the protein that transports B12 to cells. These patients may have a functional B12 deficiency even if serum levels of B12 are normal.

If levels of MMA are elevated, it may be advisable to increase B12 consumption. Common foods high in B12 include beef liver, sardines, lamb, wild caught salmon, grass-fed beef, nutritional yeast and eggs. Vitamin B12 levels can also be increased through supplementation of B12 (taken as cobalamin, methylcobalamin, hydroxycobalamin, or adenosylcobalamin). Symptoms of a vitamin B12 deficiency include: fatigue, brain fog, memory problems, muscle weakness, unsteady gait, numbness, tingling, depression, migraines/headaches and low blood pressure.

Xanthurenate

Xanthurenate (also known as xanthurenic acid) and Kynurenate (kynurenic acid) are functional markers of vitamin B6 (also known as pyridoxine) deficiency. Vitamin B6 is a critical co-factor to over 100 important reactions that occur in the human body and is stored in the highest concentrations in muscle tissue. Tryptophan is readily converted to NAD by the liver. One of the steps in this pathway requires B6. When there is insufficient B6, xanthurenate is made instead. Kynurenate may also become elevated when patients are B6 deficient because of a different, possibly less B6 dependent pathway. The pathways leading to these biomarkers have other influences, so they will not always agree. When Xanthurenate is elevated, Kynurenate is also elevated about 1/3 of the time. When both are elevated, a B6 deficiency is likely more certain and more severe. Not only is xanthurenate an indicator of a lack of B6, it is also harmful to the human body. It complexes with insulin and decreases insulin sensitivity. In fact, rats fed xanthurenate will actually develop diabetes because of the effects on insulin. If xanthurenate levels are elevated, B6 supplementation may be considered. Food high in B6 include turkey breast, grass-fed beef, pinto beans, avocado, pistachios, chicken, sesame and sunflower seeds.

While there is always some tryptophan going down the kynurenine pathway towards NAD (and possibly xanthurenate), this



| | |
|--------------------------------------|---|
| PATIENT: XXXXXXXXXXXXXXXXXXXX | TEST REF: TST-NL-XXXX |
| TEST NUMBER: T-NL-XXXXX (XXXXXXXXXX) | COLLECTED: XX/XX/XXXX |
| GENDER: XYZ | RECEIVED: XX/XX/XXXX |
| AGE: XX | TESTED: XX/XX/XXXX |
| | PRACTITIONER: XXXXXXXXXXXXXXXXXX XXXXXXXXXXXXXXXXXXXXXX |

TEST NAME: Cycle Mapping Plus

process is up-regulated by inflammation, estrogen and cortisol. If levels of estrogen or cortisol are high, it may exacerbate xanthurenate elevations and increase the need for B6. Xanthurenate can also bind to iron and create a complex that increases DNA oxidative damage resulting in higher 8-OHdG levels. If both markers are elevated, there is likely an antioxidant insufficiency.

Pyroglutamate

Pyroglutamate (also known as pyroglutamic acid) is a functional marker of glutathione deficiency. Pyroglutamate is a step in the production/recycling of glutathione. If the body cannot convert pyroglutamate forward, it will show up elevated in the urine. High pyroglutamate is an established marker for glutathione deficiency. Pyroglutamate in the urine can also be elevated with cheese consumption.

Glutathione is one of the most potent anti-oxidants in the human body. It is especially important in getting rid of toxins, including the reactive quinone species formed by 4-OH-E1 and 4-OH-E2. This reactive species can damage DNA if not detoxified by either methylation or glutathione.

Some have reported that low pyroglutamate may also be indicative of a need for glutathione; however, this is not established in the scientific literature.

Neurotransmitter Metabolites

The neurotransmitters dopamine, norepinephrine and serotonin are important for human health. Measuring neurotransmitters directly (direct testing of serotonin, for example) is difficult because of their instability and their urinary measurements are controversial with respect to how well they reflect the body's levels of these neuro-hormones. Each of these three neurotransmitters can be assessed indirectly by measuring their urine metabolites. While these metabolites are not a perfect reflection of what's going on in the brain, the scientific literature does affirm their use for a good representation of overall levels of these neurotransmitters.

Homovanillate (HVA)

Homovanillate (also known as HVA) is the primary metabolite of dopamine, a brain and adrenal neurotransmitter that comes from tyrosine (with BH4 and iron as co-factors) and goes on to create norepinephrine (noradrenaline) and epinephrine (adrenaline).

Low levels of HVA can be due to low levels of dopamine or poor conversion of dopamine to HVA. The latter may be due to insufficient levels of SAM, Magnesium, FAD and NAD which are needed to metabolize dopamine. Low circulating dopamine may be due to insufficient BH4, iron or tyrosine. It may also be seen when adrenal function is generally low. Low dopamine levels may be associated with addictions, cravings and pleasure seeking (to boost levels) in addition to sleepiness, impulsivity, tremors, less motivation, fatigue and low mood.

Elevated HVA may be caused by generally increased adrenal hormone output or because of a copper or vitamin C deficiency (which are needed for dopamine conversion to norepinephrine). Elevations may also be caused by a number of medications or supplements including: MAO inhibitors, quercetin, tyrosine, DL-phenylalanine (DLPA), L-dopa, macuna, dopamine medication (Levodopa, Sinemet, Methyldopa), SNRI medication (Wellbutrin), tricyclic antidepressants, amphetamines, appetite suppressants, and caffeine. Bananas also contain dopamine. Elevated dopamine may be associated with loss of memory, insomnia, agitation, hyperactivity, mania, hyper-focus, high stress and anxiety as well as addictions, cravings and pleasure seeking (to maintain high levels).

When HVA is very high, consider if the previously discussed foods, supplements or medications may be the cause. Rarely, tumors associated with increased HVA may be present. In these cases, further testing is necessary for diagnosis. High HVA alone is not diagnostic of a tumor.

Vanilmandelate (VMA)

Vanilmandelate (also known as VMA) is the primary metabolite of norepinephrine and epinephrine (adrenaline). The adrenal gland makes cortisol and DHEA as well as norepinephrine and epinephrine. When adrenal hormone output is generally low, VMA levels may be low. If HVA levels are significantly higher than VMA, there may be a conversion problem from dopamine to norepinephrine. This case can be caused by a copper or vitamin C deficiency. The enzymes COMT (methylation) and MAO are needed to make VMA from norepinephrine. If these enzymes are not working properly, VMA may be low when circulating norepinephrine and/or epinephrine are not low. Low levels of norepinephrine and epinephrine may be associated with addictions, cravings, fatigue, low blood pressure, low muscle tone, intolerance to exercise, depression, loss of alertness.

When the body is under physical or psychological stress, VMA levels may increase. Because dopamine gets converted to norepinephrine and ultimately to VMA, the list of medications and supplements that increase HVA may also increase VMA. Elevated levels may be associated with feeling stressed, aggression, violence, impatience, anxiety, panic, worry, insomnia, paranoia, increased tingling/burning, loss of memory, pain sensitivity, high blood pressure and heart palpitations.

If VMA and HVA are both extremely high, it may be necessary to rule out a neuroblastic tumor.

Melatonin (measured as 6-OHMS)

Melatonin is not technically an adrenal or sex hormone however it is highly involved in the entire endocrine system. It is made in small amounts in the pineal gland in response to darkness and stimulated by Melanocyte Stimulating Hormone (MSH). A low MSH is associated with insomnia, an increased perception of pain, and mold exposure. Pineal melatonin (melatonin is also made in significant quantities in the gut) is associated with the circadian rhythm of all hormones (including female hormone release). It is also made in small amounts in the bone marrow, lymphocytes, epithelial cells and mast cells. Studies have shown that a urine sample collected upon waking has levels of 6-Hydroxymelatonin-sulfate (6-OHMS) that correlate well to the total levels of melatonin in blood samples taken continuously throughout the night. The DUTCH test uses the waking sample only to test levels of melatonin production.

Low melatonin levels may be associated with insomnia, poor immune response, constipation, weight gain or increased appetite. Elevated melatonin is usually caused by ingestion of melatonin through melatonin supplementation or eating melatonin-containing foods. Elevated melatonin production that is problematic is rare, but levels can be higher in patients



| | | |
|--|------------------------------|--|
| PATIENT: XXXXXXXXXXXXXXXXXXXX | TEST REF: TST-NL-XXXX | |
| TEST NUMBER: T-NL-XXXXX (XXXXXXXXXXXX) | COLLECTED: XX/XX/XXXX | PRACTITIONER: XXXXXXXXXXXXXXXXXXXX |
| GENDER: XYZ | RECEIVED: XX/XX/XXXX | XXXXXXXXXXXXXXXXXXXX |
| AGE: XX | TESTED: XX/XX/XXXX | XXXXXXXXXXXXXXXXXXXX |

TEST NAME: Cycle Mapping Plus

with Chronic Fatigue Syndrome and may be phase shifted (peaking later) in some forms of depression.

8-OHdG (8-hydroxy-2-deoxyguanosine) results can be seen on page 6 of the DUTCH Complete (or DUTCH Plus) report. It is a marker for estimating DNA damage due to oxidative stress (ROS creation). 8-OHdG is considered pro-mutagenic as it is a biomarker for various cancer and degenerative disease initiation and promotion. It can be increased by chronic inflammation, increased cell turnover, chronic stress, hypertension, hyperglycemia/pre-diabetes/diabetes, kidney disease, IBD, chronic skin conditions (psoriasis/eczema), depression, atherosclerosis, chronic liver disease, Parkinson's (increasing levels with worsening stages), Diabetic neuropathy, COPD, bladder cancer, or insomnia. Studies have shown higher levels in patients with breast and prostate cancers. When levels are elevated it may be prudent to eliminate or reduce any causes and increase the consumption of antioxidant containing foods and/or supplements. The reference range for 8-OHdG is a more aggressive range for Functional Medicine that puts the range limit at the 80th percentile for each gender. A classic range (average plus two standard deviations) would result in a range of 0-6ng/mg for women and 0-10ng/mg for men. Seeking out the cause of oxidative stress may be more crucial if results exceed these limits.

